

Omega HW #1 – Calculating functions

Krishanu Sankar

September 21, 2025

Square roots

In class, we talked about two methods of calculating $\sqrt{2}$. In the “decimal” method, we considered the decimal representation $\sqrt{2} = 1.d_1d_2d_3d_4\dots$ and iteratively calculated d_1, d_2, d_3 , and so on. In the “fractional” method, we begin with an initial approximation x (e.g., $x = 1$), and repeatedly applied the transformation $x \mapsto \frac{1}{2} \left(x + \frac{2}{x} \right)$ to get successively better approximations.

1. (1) Calculate $\sqrt{17}$ to at least 3 decimal places, using the decimal method.
2. (2) Calculate the first three fractional approximations of $\sqrt{17}$. (*Hint: You will need to change the transformation you use.*)
3. Let $f(x) = \frac{1}{2} \left(x + \frac{2}{x} \right)$.
 - (a) (1) Calculate $f(\sqrt{2})$.
 - (b) (1) Draw a sketch of $y = f(x)$ for $1 \leq x \leq 2$. You can use a tool to help you, e.g. <https://www.desmos.com/calculator>. Indicate the following points in the graph:
 - The minimum value of the function.
 - The points $(x, f(x))$ for the first three x values in the fractional method (beginning with $x = 1$).
 - (c) (2) If x is a number in the interval $[1, 2]$, what can you say about the relative sizes of $|x - \sqrt{2}|$ and $|f(x) - \sqrt{2}|$, based on looking at this graph? Which is larger, and why?

Functions and inverses

Given a function $f(x)$, its *inverse* $f^{-1}(x)$ (if it exists!) is a function such that

- $f^{-1}(f(x)) = x$ for all x in the domain of f , and
- $f(f^{-1}(x)) = x$ for all x in the domain of f^{-1} .

1. (1) Find the inverse of $f(x) = 4 - \frac{3}{4}x$.
2. (2) Let $f(x) = 3x^2 + 4x + 3$.

- Find a function $g(x)$ such that $f(g(x)) = x$.¹
- What are the domain and range of $g(x)$?
- Draw the graphs of both $y = f(x)$ and $y = g(x)$.

Bonus question: Cube roots and beyond

- (2) Think about how you could adapt the decimal method to calculate *cube roots*. Use this method to calculate $\sqrt[3]{2}$ to 3 decimal places. I've included the first step below to get you started.²

$$\begin{array}{r}
 1. \quad 2 \quad \dots \\
 \hline
) \quad 2.000\,000\,000\dots \\
 - 1 \\
 \hline
 1 \quad 000 \\
 - 600 \\
 - 120 \\
 - 8 \\
 \hline
 272\,000
 \end{array}$$

- (3) Can you describe an algorithm to calculate $\sqrt[n]{x}$ for any whole number n ? What about $x^{m/n}$ for any whole numbers m and n ? (Remember that fractional exponents are defined by $x^{m/n} = (\sqrt[n]{x})^m$.)

¹*Hint:* Complete the square.

²Doing this by hand is computation-heavy, and requires multiplying 3-digit integers along the way. You can use a calculator to help you.